FORM B.

Return showing the description and weight of all yarn spun in the Cotton Mills in the Mysore State, during the month ending 31st August 1931.

Count of yarn	Weight in pounds of each count	Count of yarn	Weight in pounds of each count	Count of yarn	Weight in pounds of each count
2s	5,452	14s	56,375	25s	6,130
6s	34,733	14½s	34,879	26s	29,831
6½s	43,726	15s	536	27s	13,145
8s	52,366	16s	33,150	28s	67,545
8½s	34,824	16½s	3,443	30s	81,421
10s	11,170	18s	1,896	32s	12,067
10½s	15,633	19s	119,705	36s	15,987
11s	86	20s	178,332	40s	93,160
11½s	14,918	20½s	34,068	42s	834
12s	23,404	21s	54,921	60s	7
12½s	64,286	22s	29,696		
13½s	13,396	22½s	372	Total	1,262,656
		24s	81,862		

H. K. RAMIENGAR,
For *Director of Industries and Commerce, in Mysore.*

MYSORE DEPARTMENT OF HEALTH.

NOTE ON THE STATE OF PUBLIC HEALTH IN MYSORE STATE FOR THE WEEK ENDED THURSDAY, SEPTEMBER 24, 1931.

A total of 64 attacks with 55 deaths from plague was returned during the week against 140 attacks with 73 deaths in the previous week. Mysore City returned 4 attacks with 4 deaths while Bangalore City and Kolar Gold Field remain free from the disease. Comparative figures for the corresponding week in 1930 and 1929 were 96 attacks with 53 deaths and 210 attacks with 115 deaths respectively.

Two hundred and six attacks with 109 deaths from cholera were returned from one village in each of Magadi, Yedathore, Honnali, Channagiri, Tiptur and Pavagada Taluks, two villages in each of Nelamangala and Malvalli Taluks, four villages in each of Hoskote and Harihar Sub-Taluks, and in six villages in each of Jagalur and Davangere Taluks. Ajjampur and Tiptur Towns including Bangalore City reported cases.

One fatal case from influenza was reported from Mysore City.

The State remained free from small-pox and Relapsing Fever.

The total mortality for Bangalore City amounted to 92, representing a calculated annual ratio of 40·35 per mille of population against the average ratio of 43·98 for the corresponding week of last year. Classified according to causes, the deaths numbered 2 under cholera, 2 under malaria, 2 under broncho-pneumonia, 10 under pneumonia, 2 under typhoid, 1 under other fevers, 4 under phthisis, 4 under dysentery and diarrhœa and 65 under all other causes.

During the week, the Civil and Military Station, Bangalore, returned 79 deaths representing a calculated annual ratio of 34·45 per mille of population against the average ratio of 39·28 for the corresponding week of last year. Classified according to causes, the deaths were—broncho-pneumonia 10, pneumonia 5, other fevers 3, phthisis 6, dysentery and diarrhœa 5, and all other causes 50.

The total mortality for Mysore City was 61 representing a calculated annual ratio of 37·78 per mille of population against the average ratio of 39·03 for the corresponding week of last year. Classified according to causes, the deaths were—plague 4, malaria 2, broncho-pneumonia 1, pneumonia 7, typhoid 2, phthisis 3, dysentery and diarrhœa 11, and all other causes 31.

There were 37 deaths from all causes in Kolar Gold Field equivalent to a calculated annual ratio of 21·95 per mille of population against the average ratio of 29·76 for the corresponding week of last year. Classified according to causes, the deaths numbered, 2 under pneumonia, 4 under other fevers, 2 under phthisis, 8 under dysentery and diarrhœa, and 21 under all other causes.

J. V. KARVE,
For *Director of Health.*